

The good news is that ADHD can be diagnosed and treated at any age. Adults can benefit from treatment, including medication and therapy, as much as children and adolescents. If you are an adult and have concerns that you might have ADHD, it certainly is worthwhile to consider pursuing a diagnostic evaluation. The risks associated with undiagnosed and untreated ADHD in adulthood are many, including increased risk for job and career failure, marital problems and divorce, financial problems and substance abuse.

The Comprehensive Evaluation for ADHD

A diagnostic evaluation can be provided by a qualified licensed health care professional. It should be said that ADHD cannot be diagnosed accurately from just brief office observations or from talking to the person. Clinicians will vary somewhat in the procedures and testing materials they use; however, certain protocols are considered essential for a comprehensive evaluation. A comprehensive diagnostic assessment should include:

- A thorough diagnostic interview, including a detailed history of past and current functioning
- A comprehensive history (developmental, medical, academic, work, social, family)
- ADHD symptom checklists
- Standardized behavior rating scales for ADHD
- Other types of psychometric testing as deemed necessary by the clinician
- Review of past evaluations and school records
- Screening for the presence of other co-morbid conditions

The Structured Clinical Interview and Review of Records

The single most important part of a comprehensive evaluation for ADHD is a structured or semi-structured interview to provide a detailed history of past and current behavior for the individual. This interview allows the clinician to cover a broad range of topics, discuss relevant issues in more detail and ask follow up questions. The examiner will likely ask questions about the person's health history, developmental history going back to early childhood, academic history, work history, family and marital history and social history.

Many professionals find it helpful to review old report cards and other school records, going back to kindergarten or preschool. If such records are available, they should be brought to the first appointment. Copies of reports from any previous psychological testing should also be brought to the appointment. For adults who experience problems in the workplace, job evaluations should be brought for review if these are available.

Participation of a Significant Other

When possible, it is important for the clinician to interview a significant other (spouse, other family member, parent, etc.) who knows the person well. Many adults with ADHD report having a spotty or poor memory of their past, particularly memories of their childhood. They may recall specific details, but forget important events or other facts. The person being evaluated may be asked to have his or her parents or an adult sibling fill out a retrospective ADHD questionnaire describing childhood behavior.

Gender Difference Considerations

Clinical guidelines for diagnosis of ADHD are provided in the diagnostic manual of the American Psychiatric Association, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, also known as the DSM-IV (American Psychiatric Association, 2000). For a variety of reasons, ADHD appears to be under-diagnosed in females. DSM-IV diagnostic criteria are not as well suited for diagnosing ADHD in girls and women. Girls with ADHD are less likely to be hyperactive compared to boys their age, and consequently are less disruptive and less apt to draw attention from the adult caretakers in their lives. The onset of ADHD symptoms generally occurs later in girls, frequently not before the middle school years. Women with ADHD are more likely to experience co-existing depression and anxiety disorders. These gender differences should be considered when reviewing the history and assessing current symptoms in female patients.

ADHD Symptom Checklists and Standardized Behavior Rating Scales

During an evaluation, the clinician uses the DSM-IV to determine the extent to which the symptoms found in the diagnostic criteria apply to the individual currently and since childhood. Other self-report and collateral report standardized behavior-rating scales may be used, such as the Brown ADD Scales-Adult Version and the Conners Adult Attention Rating Scale.

Limitations of DSM-IV Diagnostic Criteria

The DSM–IV diagnostic criteria for ADHD were developed based on data from children between the ages of four and 17 (Brown, 2000). Consequently, it appears that these criteria are ideally suited for the diagnosis of children (particularly boys) with ADHD. Unfortunately these existing criteria are not well suited for diagnosis of ADHD in adults, and likely underestimate the true rate of ADHD in the adult population.

The DSM–IV criteria do not adequately assess cognitive impairments associated with ADHD, particularly impairments in executive functioning. Although the DSM–IV requires onset of some symptoms before age seven, onset of symptoms may not be evident in some individuals until well past that age. Impairment may not be evident in individuals with high intelligence and good coping skills until the person is in high school, in college, in a demanding career position, or in the demanding role of parent and household manager.

Until improved diagnostic criteria for ADHD in adults are made available, clinicians are advised to use the existing DSM–IV criteria for guidance in diagnosis of adults. Clinicians are encouraged to be aware of individual differences in onset and impact of ADHD symptoms and to make an appropriate diagnosis on the basis of good clinical judgment even when the DSM-IV criteria are not strictly met for adult patients.

Tests of Vigilance and Attention

Some clinicians prefer to use computerized tests of attention as a source of additional information. These may involve, for example, pressing a key when a letter or symbol appears on the screen. The computerized tests are not diagnostic by themselves, but are used as a supplement and can provide additional information useful in making a diagnosis. Many individuals with ADHD perform well on these tests, therefore, it should be noted that obtaining a “normal” score does not rule out having ADHD.

The Role of Psychological Testing in Adult ADHD Evaluations

Depending on the individual and the concerns being addressed, additional psychological, neuropsychological or psychoeducational testing may be used as needed. These tests do not diagnose ADHD directly and should not be used by themselves to evaluate for ADHD. Such tests can provide important information about ways in which ADHD impacts various areas of functioning for the individual, notably emotional and intellectual functioning. In particular,

psychological testing can help determine the presence and effects of co-existing conditions.

Screening and Assessment of Co-Existing Conditions

Research has shown that 60 percent or more of individuals with ADHD have one or more co-existing conditions. The most common include depression, anxiety disorders, bipolar disorder, substance abuse and addictions and learning disabilities. Many coexisting conditions mimic ADHD symptoms (distractibility, restlessness, forgetfulness, etc.) and may be mistaken for the disorder. When there is one or more co-existing conditions with ADHD, it is essential that all are diagnosed and treated. Failure to treat co-existing conditions often leads to failure in treating the ADHD. If there are indications of substance abuse, either currently or in the past, a detailed history of substance use and abuse should be taken.

The Medical Examination

Some medical conditions (for example, thyroid problems or seizure disorders) may cause symptoms that mimic ADHD or may co-exist and complicate the condition. If the individual being evaluated for ADHD has not had a recent physical exam (within 6–12 months), a medical examination is recommended to rule out medical causes for symptoms. The examining physician may also consider laboratory testing to screen for medical conditions if specific symptoms are uncovered during the interview.

Providing Feedback and Treatment Recommendations

The clinician should integrate the information that has been collected from interviews and testing forms, complete a written summary or report, and provide the person with diagnostic opinions concerning ADHD as well as any other co-existing conditions. The clinician should review treatment options and assist the individual in planning a course of appropriate medical and psychosocial treatment intervention. Afterwards, the clinician should communicate with the individual's primary care providers, as deemed necessary.

Peter Jaksa, PhD, is a licensed clinical psychologist in Chicago, Ill., and clinical director of ADD Centers of America LLC (www.addcenters.com). He is the parenting editor for ADDitude Magazine and serves on the Scientific Advisors Board. He is a past president of the Attention Deficit Disorder Association.